



SCGC Service Description

Updated December 14, 2021

To complete an SCGC service inquiry or request, start [here](#).

Table 1. List of SCGC services. See detailed description below.

Service	Cat #	Unit	Unit price*
SAG Generation			
SAG Generation 2	S-201	384-well plate	\$3,800
SAG SEQUENCING AND BIOINFORMATICS			
SAG WGS 1.1 billion reads	S-211	≤384 SAGs	\$24,000
SAG WGS 0.4 billion reads	S-212	≤384 SAGs	\$19,000
SAG WGS 0.1 billion reads	S-216	≤384 SAGs	\$15,000
SAG WGS addon 1.1 billion reads	S-213	≤384 SAGs	\$14,000
SAG WGS addon 0.4 billion reads	S-214	≤384 SAGs	\$9,000
SAG WGS addon 0.1 billion reads	S-215	≤384 SAGs	\$6,000
ADDITIONAL SERVICES			
SAG re-arraying	S-105	96-well plate	\$450
Sample cryoprotectant glyTE	S-019	10 mL	\$100
Consultation	S-011	1 hour	\$460
Customized Services	S-100	Custom	Request a quote

*All prices are in US dollars (USD). 15% discount is applied if SCGC services that exceed \$200,000 in value are ordered simultaneously. 20% discount is applied on all services for Bigelow Laboratory's in-house projects. Service fees do not include shipping costs.



OVERVIEW

Single cell DNA sequencing

Single cell genomics unveils the genomic blueprints of the most fundamental units of life. It is a powerful approach to analyze biochemical properties, evolutionary histories and the biotechnological potential of uncultured microorganisms, which constitute over 99% of biological diversity on our planet. Single cell genomics is also emerging as a revolutionary technology in the studies of cancer, autoimmune diseases and hereditary disorders. Single cell genomics consists of a series of integrated processes, starting with appropriate sample collection and preservation, followed by physical separation, lysis and whole genome amplification of individual cells, then proceeding to DNA sequencing and sequence interpretation (1). These processes are incorporated in the comprehensive suite of services offered by SCGC (Table 1).

The Single Cell Genomic Center (SCGC®) at Bigelow Laboratory for Ocean Sciences

The SCGC is world's first research and service center with the primary focus on the single cell genomics of microorganisms (see About SCGC). This includes microorganisms from diverse microbiomes, biosafety level 2 organisms and organisms from hard-to-process environments, such as soil and the deep biosphere. SCGC also has experience and capabilities to process individual cells of humans and other multicellular organisms.

Commitment to Quality

The processing of exceedingly small DNA quantities makes single cell genomics highly susceptible to DNA contamination and amplification biases. At SCGC, we take these risks very seriously and have developed techniques to minimize and monitor methodological artifacts at each point in our workflow. Single cell/particle sorting and DNA amplification are performed in a cleanroom environment, and all associated consumables are decontaminated using methods that were developed and evaluated at SCGC (2). Single amplified genome (SAG) generation services include multiple controls to detect potential DNA contamination. To prevent index switching during SAG sequencing, multiplexed libraries contain dual, 10 bp barcodes. Prior to *de novo* assembly, raw reads are quality-filtered and digitally normalized using in-house, optimized protocols (2). Genome *de novo* assemblies are analyzed using multiple QC algorithms. The entire workflow is evaluated for contamination and assembly errors using microbial benchmark cultures with diverse genome complexity and GC content (%), indicating no non-target and undefined bases and average frequencies of mis-assemblies, indels and mismatches at <5 per 100 kbp (2).



SCGC SERVICES: SAG GENERATION

Single amplified genomes (SAGs) are products of whole genome amplification reactions performed on individual cells or other DNA-containing particles (1, 3). SAG generation involves physical separation, lysis and DNA amplification of individual cells/particles. SCGC has developed state-of-the-art techniques for each of these steps (2) and combined them in service S-201.

SAG-Generation (S-201)

The S-201 service includes separation of individual cells or DNA-containing particles into wells of a 384-well plate by fluorescence-activated cell sorting (FACS), followed by cell lysis and genomic DNA amplification. Cells/particles are separated using an inFLux Mariner (BD), which can be finely tuned to select individual cells or particles based on a range of optical characteristics. Cells/particles may be selected based on the particle autofluorescence, the fluorescent DNA stain SYTO-9 (Thermo Fisher Scientific, provided by SCGC) or other stains and probes that are applied by an SCGC customer prior to shipping to SCGC. During sorting, multiple wells are used as negative and positive controls on each plate (Figure 1).

Cells/particles are lysed and their DNA is denatured by 2 freeze-thaw cycles and a subsequent KOH treatment (2). SCGC uses WGA-X[®] for genomic DNA amplification, a method developed by SCGC (2). Compared to the earlier versions of multiple displacement amplification technique (MDA) (4), WGA-X improves average genome recovery from individual cells and viral particles, with most notable enhancements observed in SAGs with high G+C content (Figures 2 and 3). Please note that SAG generation success varies among samples (Figure 1) and depends on many factors, such as: a) prompt cryopreservation of intact cells and gDNA prior to cell sorting; b) successful discrimination of cells and viral particles from other particles during FACS; c) successful single cell and viral particle lysis and DNA amplification. Deliverables of S-201 include:

- a) One 384-well microplate containing WGA-X products of individual cells or DNA-containing particles, 10 uL per well, usually averaging ~1 microgram gDNA per well.
- b) Index FACS data files.
- c) WGA-X kinetics data obtained by measuring fluorescence of a DNA stain in each well.



For a standard S-201 service fee to be applicable, the following conditions must be met:

- a) Target cells or particles are in aquatic solution, are less than 40 micrometers in diameter, and are cryopreserved and shipped following SCGC recommendations (see Preparation and Shipment on SCGC website). Samples that have not been brought into aquatic suspension by SCGC customers may be analyzed as a customized service. Please contact SCGC manager Brian Thompson for additional advice on how to produce SCGC-compatible samples in your own lab or for a feasibility assessment and a quote.
- b) Sorting is based on particle autofluorescence, SYTO-9 fluorescence (DNA stain provided by SCGC), or probes that are applied by SCGC customer prior to shipping to SCGC.
- c) Only one sample is used per microplate, following SCGC's standard plate setup (Figure 1).




Figure 1. Examples of WGA-X kinetics in microplates with divergent DNA amplification success rates. The reaction Cp value indicates time (hours) required to reach half of the maximal DNA-SYTO-9 fluorescence. The Cp is inversely correlated with amplification speed and the initial quantity of gDNA that is available to polymerase.

b) -




Figure 2. Correlation between whole genome amplification Cp and genome recovery from SAGs of benchmark strains. Each bacterial strain data set derives from eight randomly selected SAGs and five SAGs with the lowest Cp on a 384-well plate. Each eukaryote data set derives from three SAGs with the lowest Cp. Assemblies of bacterial and eukaryote SAGs were produced from five million and twenty millions of 2×150 bp reads, respectively. From (2).




Figure 3. Genome recovery from SAGs of benchmark microorganisms as a function of sequencing effort. Paired-end reads of each bacterial SAG were randomly down-sampled to 0.1, 0.5, 1, 2, 3, 4, and 5 millions. Paired-end reads of each eukaryote SAG were randomly down-sampled to 5, 10 and 20 millions. Note that data markers are slightly offset on the horizontal scale, in order to better discriminate standard deviation bars among the treatments. The inset provides a better view of results at low read numbers. Displayed are averages and standard deviations, which were estimated from eight randomly selected SAGs in the case of bacterial SAGs, and from three SAGs with the lowest reaction Cp values in the case of eukaryote SAGs. From (2).

SCGC SERVICES: SAG SEQUENCING AND BIOINFORMATICS

SCGC's SAG whole genome sequencing (WGS) services combine library preparation, shotgun sequencing, *de novo* assembly, annotation and quality control. Each of these steps are optimized for single cell genomics and validated using benchmark cultures (2). Please note that SCGC does not process SAGs that have been handled elsewhere, in order to prevent the risk of DNA contamination at our facility.

Sequencing libraries are prepared with Nextera XT (Illumina) reagents using a modified protocol (2). Multiplexed libraries are sequenced with Illumina's NextSeq 2000. SCGC's *de novo* genome assembly workflow involves quality-trimming of raw reads with Trimmomatic (5), removal of low-complexity and reagent contaminant reads, read normalization with kmernorm, assembly with SPAdes (6), quality-trimming of contigs, and the removal of contigs shorter than 2000 bp, as previously described (2). Functional annotation is performed using



Prokka (7), complemented with a custom protein annotation database built from compiling Swiss-Prot (8) entries for Archaea and Bacteria. SAG taxonomic assignments are based on SSU rRNA gene phylogeny and checkM (9). Estimates of genome assembly completion and potential contamination are obtained with checkM. A complementary search for contaminant sequences is performed using tetramer principal component analysis (10). Please note that our current annotation tools are designed for bacterial and archaeal genomes and are not suitable for eukaryotes. Deliverables of all SCGC WGS services include:

- a) raw sequence reads
- b) *de novo* SAG assemblies
- c) functional annotation of bacterial and archaeal SAGs
- d) SAG taxonomic assignments
- e) general genome properties, such as GC content and coding density
- f) QC analyses

Specialized laboratory and computational procedures are necessary in order to obtain high quality genomic sequences from SAGs. A particular challenge to consider is the cross-talk of multiplexed libraries (also known as “sample bleeding” and “index switching”) (11, 12). *De novo* assemblies of SAGs are particularly susceptible to this problem, due to the fact that single cell whole genome amplification is highly uneven across the genome, and deep sequencing is typically employed to facilitate the recovery of the under-amplified genome regions (1). As a result, even a relatively small overall fraction of miss-assigned reads may form contigs that represent over-amplified regions of co-sequenced SAGs. SCGC uses in-house infrastructure and procedures for SAG sequencing that eliminate library cross-talk and do not require subsequent, computational decontamination (2). To verify the efficacy of these solutions, SCGC benchmarks its entire workflow using SAGs of previously sequenced strains (2). We encourage SCGC customers to take similar precautions and workflow validation when sequencing SAGs outside SCGC, in order to maintain data integrity.

SAG WGS 1.1 billion reads (S-211)

This is the most cost-effective option for sequencing of large batches of SAGs. DNA sequencing is performed using an Illumina NextSeq2000 instrument and P3 reagents in 2×100 bp mode, resulting in a total of ~1.1 billion paired-end reads. Raw reads are processed using our standard bioinformatics assembly, annotation and QC workflow, as described above. A total of up to 384 SAGs can be selected for sequencing from up to six, 384-well SAG plates generated using SCGC services S-201. SAG selection can be done by the customer or by SCGC,



e.g.; at random or based on the WGA-X Cp values, which correlate with SAG's potential for high genome recovery (Figure 2). The average SAG sequencing depth depends on the number of SAGs selected by a customer for sequencing, which can be guided by our prior results with benchmark cultures (Figure 3). Some SAGs may generate low read numbers, due to physical, biological or biochemical factors of individual cells or samples, or due to liquid handling inconsistencies. Due to the high-throughput, low-cost nature of SCGC WGS services, they do not include free re-sequencing of such SAGs, unless the number of affected SAGs exceeds 10% of all sequenced SAGs.

SAG WGS 0.4 billion reads (S-212)

This service is similar to S-211; the only differences are: a) use of P2 sequencing reagents; b) lower cost; and c) the total read count is ~0.4 billion, i.e.; ~3x lower than with service S-211. Service S-212 is best suited for full-depth sequencing of up to 100 prokaryote SAGs or for low-coverage sequencing of up to 384 SAGs.

SAG WGS 0.1 billion reads (S-216)

This service is similar to S-211; the only differences are: a) use of P1 sequencing reagents; b) lower cost; and c) the total read count is ~0.1 billion, i.e.; ~11x lower than with service S-211. Service S-216 is best suited for full-depth sequencing of up to 30 prokaryote SAGs or for low-coverage sequencing of up to 384 SAGs.

SAG WGS addon 1.1 billion reads (S-213)

This service includes deeper sequencing, *de novo* assembly, functional annotation and QC of up to 384 SAGs for which compatible Illumina libraries have already been produced by a single service S-211, S-212 or S-216. The total number of new, 2x100 bp reads produced by this service is ~1.1 billion and is distributed relatively evenly among the selected SAGs. More reads per SAG usually results in better genome recovery, but this relationship is not linear (Figure 3). Per customer's request, SCGC can generate lists of SAGs that meet S-213 eligibility criteria and are either randomized or prioritize SAGs with lowest WGA Cp values – indicators of the potential for high genome recovery (Figure 2). Please note that, due to potential liquid handling inconsistencies, some SAGs may receive unexpectedly low numbers of sequence reads. Due to its high-throughput, low-cost nature, service S-213 does not include free re-sequencing of such SAGs, unless the number of affected SAGs exceeds 10% of all sequenced SAGs.

**SAG WGS addon 0.4 billion reads (S-214)**

This service is similar to S-213. The only differences are: a) use of P2 sequencing reagents; b) lower cost; and c) the total read count is ~0.4 billion, i.e.; ~3x lower than with service S-213.

SAG WGS addon 0.1 billion reads (S-215)

This service is similar to S-213. The only differences are: a) use of P1 sequencing reagents; b) lower cost; and c) the total read count is ~0.1 billion, i.e.; ~11x lower than with service S-213.

SCGC SERVICES: ADDITIONAL SERVICES**SAG re-arraying (S-105)**

SAGs are transferred from the original 384-well plates to 96-well plates. The SCGC customer defines transfer volumes, source wells, and destination wells. Prior to the transfer, the destination wells are pre-filled with 5-150 uL of either deionized water or 1x TE buffer, as specified by the SCGC customer. Deliverables include re-arrayed SAGs in a 96-well plate.

Sample cryoprotectant glyTE (S-019)

For SCGC customers' convenience, we offer the preparation and shipment of SCGC's recommended sample cryoprotectant glyTE. Deliverables of this service include preparation and shipment of 10 mL of 10x concentrated glyTE. Please note that SCGC's recommended procedures for sample cryopreservation, including the recipe for the cryoprotectant glyTE, are posted on the SCGC website under Preparation and Shipment.

Consultation (S-011)

Basic support is included in the pricing of all SCGC services. When SCGC customers require more extensive help with study design and/or data interpretation (greater than two hours per project), we may request compensation for the associated labor costs via collaborative research grants or consultant fees. We may also request co-authorship on resulting publications for



those SCGC scientists who are providing substantial, project-specific intellectual input.

Customized services (S-100)

SCGC offers a wide range of customized services, e.g., non-standard cell sorting, non-SAG sequencing, bioinformatics support, method development, etc. Co-authorship on resulting publications may be requested for those SCGC scientists who are providing substantial, project-specific intellectual input. For more information, please contact SCGC manager Brian Thompson.

COST ESTIMATE: CASE STUDY

Let's assume an external SCGC customer wants to a) generate two 384-well plates of prokaryote single amplified genomes (SAGs); b) perform whole genome sequencing of 150 best-quality SAGs from each of the two plates, as defined by their lowest WGA-X Cp (Figure 2). Fees for this work are summarized in Table 2. Deliverables include:

1. Two 384-well microplates containing SAGs with associated indexFACS and gDNA amplification kinetics data.
2. 300 assembled, annotated and QC-ed SAG genome sequences.

Table 2. Cost estimate for a hypothetical SCGC project.

Service	Cat. #	Price per unit	# of units	Amount
SAG Generation 2	S-201	\$3,800	2	\$7,600
SAG WGS 1.1 billion reads	S-211	\$24,000	1	\$24,000
Total				\$33,600

REFERENCES

1. Stepanauskas R. 2012. Single cell genomics: An individual look at microbes. *Current Opinion in Microbiology* 15:613-620.



2. Stepanauskas R, Fergusson EA, Brown J, Poulton NJ, Tupper B, Labonté JM, Becraft ED, Brown JM, Pachiadaki MG, Povilaitis T, Thompson BP, Mascena CJ, Bellows WK, Lubys A. 2017. Improved genome recovery and integrated cell-size analyses of individual uncultured microbial cells and viral particles. *Nat Commun* 8:84.
3. Stepanauskas R, Sieracki ME. 2007. Matching phylogeny and metabolism in the uncultured marine bacteria, one cell at a time. *Proceedings of the National Academy of Sciences of the United States of America* 104:9052-9057.
4. Dean FB, Hosono S, Fang LH, Wu XH, Faruqi AF, Bray-Ward P, Sun ZY, Zong QL, Du YF, Du J, Driscoll M, Song WM, Kingsmore SF, Egholm M, Lasken RS. 2002. Comprehensive human genome amplification using multiple displacement amplification. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 99:5261-5266.
5. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114-2120.
6. Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Pribelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *Journal of Computational Biology* 20:714-737.
7. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068-2069.
8. Bateman A, Martin MJ, O'Donovan C, Magrane M, Alpi E, Antunes R, Bely B, Bingley M, Bonilla C, Britto R, Bursteinas B, Bye-Ajee H, Cowley A, Da Silva A, De Giorgi M, Dogan T, Fazzini F, Castro LG, Figueira L, Garmiri P, Georghiou G, Gonzalez D, Hatton-Ellis E, Li W, Liu W, Lopez R, Luo J, Lussi Y, MacDougall A, Nightingale A, Palka B, Pichler K, Poggioli D, Pundir S, Pureza L, Qi G, Rosanoff S, Saidi R, Sawford T, Shypitsyna A, Speretta E, Turner E, Tyagi N, Volynkin V, Wardell T, Warner K, Watkins X, Zaru R, Zellner H, Xenarios I, Bougueleret L, Bridge A, Poux S, Redaschi N, Aimo L, ArgoudPuy G, Auchincloss A, Axelsen K, Bansal P, Baratin D, Blatter MC, Boeckmann B, Bolleman J, Boutet E, Breuza L, Casal-Casas C, De Castro E, Coudert E, Cuche B,



Doche M, Dornevil D, Duvaud S, Estreicher A, Famiglietti L, Feuermann M, Gasteiger E, Gehant S, Gerritsen V, Gos A, Gruaz-Gumowski N, Hinz U, Hulo C, Jungo F, Keller G, Lara V, Lemercier P, Lieberherr D, Lombardot T, Martin X, Masson P, Morgat A, Neto T, Nouspikel N, Paesano S, Pedruzzi I, Pilbout S, Pozzato M, Pruess M, Rivoire C, Roechert B, Schneider M, Sigrist C, Sonesson K, Staehli S, Stutz A, Sundaram S, Tognolli M, Verbregue L, Veuthey AL, Wu CH, Arighi CN, Arminski L, Chen C, Chen Y, Garavelli JS, Huang H, Laiho K, McGarvey P, Natale DA, Ross K, Vinayaka CR, Wang Q, Wang Y, Yeh LS, Zhang J. 2017. UniProt: The universal protein knowledgebase. *Nucleic Acids Research* 45:D158-D169.

9. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *PeerJ PrePrints* 2:e1346.
10. Woyke T, Xie G, Copeland A, González JM, Han C, Kiss H, Saw JH, Senin P, Yang C, Chatterji S, Cheng JF, Eisen JA, Sieracki ME, Stepanauskas R. 2009. Assembling the marine metagenome, one cell at a time. *PLoS ONE* 4.
11. Mitra A, Skrzypczak M, Ginalski K, Rowicka M. 2015. Strategies for achieving high sequencing accuracy for low diversity samples and avoiding sample bleeding using Illumina platform. *PLoS ONE* 10.
12. Sinha R, Stanley G, Gulati GS, Ezran C, Travaglini KJ, Wei E, Chan CKF, Nabhan AN, Su T, Morganti RM, Conley SD, Chaib H, Red-Horse K, Longaker MT, Snyder MP, Krasnow MA, Weissman IL. 2017. Index switching causes “spreading-of-signal” among multiplexed samples in Illumina HiSeq 4000 DNA sequencing. *bioRxiv*:125724.